

Orientational effect of uniaxially deformed aerogel on the order parameter of superfluid 3He

E.V. Surovtsev

*P.L.Kapitza Institute for Physical Problems,
ul. Kosygina, 2, Moscow 119334, Russia*

Abstract

Effect of uniaxially-deformed aerogel on orientation of the order parameter of superfluid 3He is considered. Approach used [1] takes into account correlations in positions of particles forming aerogel. Calculations show that for the case of the B-like phase in the uniaxially-stretched aerogel deformation on the order of $10^{-2} \div 10^{-4}$ is already strong enough to achieve orientation of the order parameter which is different from that required by magnetic field in bulk liquid. For the A-like phase in uniaxially-squeezed aerogel the order of magnitude of the orientational effect is estimated. It is shown that the correlations in positions of particles forming aerogel has stronger effect on the orientation of the order parameter than on the transition temperature.

Superfluid 3He is a convenient system for investigation of the effect of impurities on formation and orientation of the order parameter. Interaction between aerogel and the order parameter of 3He can be revealed in suppression of the superfluid transition temperature, as well as in possible changing of a form of the order parameter and spatial orientation of the order parameter. The first attempts to evaluate the effect of the anisotropy of aerogel on the orientation of the order parameter were based on the Homogeneous Scattering Model (HSM) Ref.[2], which is a generalization of the Abrikosov-Gor'kov theory of superconducting alloys for the case of p-wave pairing. But this theory does not take into account correlations in positions of particles forming aerogel. Recently I.A. Fomin has shown, that such correlations can play an important role effecting on the transition temperature to the superfluid state Ref.[1]. In the present paper the effect of uniaxially deformed

aerogel on the orientation of the order parameter is considered within the same approach.

In what follows we restrict our consideration to a temperature region in the vicinity of the bulk transition temperature T_b . In this region for our purposes one can use linearized Ginzburg and Landau equation. Linearized Ginzburg and Landau equation in momentum representation in the notations of Ref.[1] can be written in the following form:

$$\left(\tau\delta_{jl} - \frac{3}{5}\xi_s^2 k^2(\delta_{jl} + 2k_j k_l)\right) a_{\mu l}(\mathbf{k}) = \int a_{\mu l}(\mathbf{q}) \eta_{jl}(\mathbf{k} - \mathbf{q}) \frac{d^3 q}{(2\pi)^3}, \quad (1)$$

$$A_{\mu j}(\mathbf{r}) = \int a_{\mu j}(\mathbf{q}) \exp(-i\mathbf{q}\mathbf{r}) \frac{d^3 q}{(2\pi)^3}, \quad \eta_{jl}(\mathbf{r}) = \int \eta_{jl}(\mathbf{q}) \frac{d^3 q}{(2\pi)^3}$$

where $\tau = \frac{T-T_b}{T_b}$, $A_{\mu j}$ - 3×3 complex matrix, corresponding to the order parameter in the case of p-wave pairing, $\eta_{jl}(\mathbf{r})$ - real symmetric tensor, which characterizes interaction between impurities and the order parameter. The form of the tensor $\eta_{jl}(\mathbf{r})$ depends on a structure of aerogel and a type of quasiparticle scattering by aerogel particles. We assume here, that aerogel consists of spheres with uniform radii ρ , distributed with an averaged density n . For simplicity the type of scattering is chosen to be diffusive. Assuming low concentration of impurities and using the Rainer and Vuorio theory of small objects in superfluid ^3He one can find the tensor $\eta_{jl}(\mathbf{r})$ to be:

$$\eta_{jl}(\mathbf{r}) = \sum_s \eta_{jl}^1(\mathbf{r} - \mathbf{r}_s), \quad (2)$$

where s - number of impurity,

$$\eta_{jl}^1(\mathbf{r}) = -\frac{\rho^2}{r^2} \nu_j \nu_l \ln \left[\tanh \left(\frac{r}{2\xi_0} \right) \right]. \quad (3)$$

As it was shown in Ref.[1], for the natural parameters of aerogel and superfluid ^3He the rhs of (1) can be treated as perturbation.

In order to find orientational correction to the free-energy one has to solve the problem of eigenvalues of Eq. (1). In the absence of perturbation the eigenvalues of Eq. (1) are degenerate, i.e single temperature transition corresponds to different orbital components of the order parameter (different orientations of the order parameter). Nominally isotropic aerogel produces isotropic perturbation. This situation was considered in Ref.[1]. In the case of such isotropic perturbation the eigenvalues of Ginzburg and Landau equation remain degenerate. In the presence of anisotropic perturbation induced by uniaxially deformed aerogel this degeneracy is partly lifted. Solution of




Figure 1:

the secular equation of perturbation theory yields corrections to the transition temperature, which correspond to "longitudinal" (along the direction of deformation) and "transverse" transition temperatures. Therefore a term with a tensor of transition temperature appears in the free-energy. The diagonal terms of this tensor are given by solution of the secular equation. After separation of isotropic and anisotropic contributions to the transition temperature orientational term in the free-energy can be expressed in a form:

$$\tau_{jl}^a A_{\mu j} A_{\mu l}^*, \quad (4)$$

$\tau_{ll}^a = 0$. Below the method of Green's functions is used.

The Green's function tensor of Eq. (1), averaged over all realizations of the potential $\eta_{jl}(\mathbf{r})$, has the form:

$$\langle G_{jl}(\tau, \mathbf{k}, \mathbf{k}') \rangle = (2\pi)^3 \delta(\mathbf{k} - \mathbf{k}') \left[\left(G_{jl}^{(0)}(\tau, \mathbf{k}) \right)^{-1} - \langle \Sigma_{jl}(\tau, \mathbf{k}) \rangle \right]^{-1}, \quad (5)$$

where

$$\left(G_{jl}^{(0)}(\tau, \mathbf{k}) \right)^{-1} = \tau \delta_{jl} - \frac{3}{5} \xi_s^2 k^2 (\delta_{jl} + 2k_j k_l), \quad (6)$$

is the unperturbed Green's function tensor and the tensor of the Self-energy part is shown diagrammatically on the Fig.1. Arrows on the figure represent the unperturbed Green's function tensor, and wavy lines correspond to the impurity potential:

$$\eta_{jl}(\mathbf{k} - \mathbf{k}') = \eta_{jl}^1(\mathbf{k} - \mathbf{k}') \sum_s e^{i(\mathbf{k} - \mathbf{k}') \cdot \mathbf{r}_s}. \quad (7)$$

Integration is made over momenta of internal lines. It is significant that the expression (5) differs from the corresponding expression of Ref.[1] in the way that it is not proportional to δ_{jl} , but it has more complicated tensor structure.

The eigenvalues of Eq. (1) are determined by the poles of Green's function, which are found from the secular equation:

$$\det \left(\delta_{jl} \cdot \tau - \frac{3}{5} \xi_s^2 k^2 (\delta_{jl} + 2k_j k_l) - \Sigma_{jl}(\mathbf{k}) \right)_{\mathbf{k} \rightarrow 0} = 0. \quad (8)$$

Here the limit $k \rightarrow 0$ corresponds to the "mobility edge" Ref. [1].

The first order correction to the Self-energy part comes from the first term of series Fig.1. Averaging over coordinates of particles renders:

$$\tau_{ba}^{(1)} = \frac{1}{3} \eta_{ll}^{(1)}(\mathbf{k} \rightarrow 0) n = \frac{\pi^2}{4} \frac{\xi_0}{l_{tr}}, \quad (9)$$

where l_{tr} - mean free path, ξ_0 - coherence length of superfluid 3He .

The effect that we are interested in appears starting from the second order correction:

$$\Sigma_{jl}^{(2)}(\mathbf{k}) = \int \eta_{jm}^{(1)}(\mathbf{k} - \mathbf{k}_1) \eta_{ln}^{(1)}(\mathbf{k}_1 - \mathbf{k}) n \langle \sum_t e^{i(\mathbf{k}_1 - \mathbf{k}) \mathbf{r}_{st}} \rangle G_{mn}^{(0)}(\tau = 0, \mathbf{k}_1) \frac{d^3 k_1}{(2\pi)^3}, \quad (10)$$

where averaged sum

$$\langle \sum_t e^{i(\mathbf{k}_1 - \mathbf{k}) \mathbf{r}_{st}} \rangle = S(\mathbf{k}_1 - \mathbf{k}), \quad (11)$$

equals by definition to the structure factor of aerogel, which characterizes correlations in positions of particles forming aerogel. The structure factor can be expressed in terms of the pair correlation function via Fourier transformation:

$$S(\mathbf{k}) = n \int C(\mathbf{r}) e^{i\mathbf{k}\mathbf{r}} d^3 r. \quad (12)$$

In accordance with the general rules of perturbation theory one substitutes in (10) the unperturbed Green's function, what is made by implying the condition $\tau = 0$.

Deformation of aerogel changes its structure factor. It is assumed here that non-deformed aerogel is isotropic. As a result its structure factor does not depend on the direction of wave-vector $\mathbf{k}$. Aerogel has several typical lengthscales. On the scales greater than several thousands $\text{\AA}$ aerogel is homogeneous. There is also an interval of scales where aerogel reveals its fractal structure. It means, that in the corresponding interval of distances $\rho < r < R$ its pair correlation function has fractal behavior:

$$C(\mathbf{r}) = A(R/r)^{3-D}, \quad (13)$$

where D - the fractal dimension of aerogel, R - the correlation radius of aerogel, i.e. distance whereon correlation between particles decreases. In order to provide smooth decreasing of correlation function on the distances of correlation radius $C(\mathbf{r})$ is replaced by the following model correlation function:

$$C(\mathbf{r}) \longrightarrow C(\mathbf{r}) \exp(-r/R). \quad (14)$$




Figure 2:

Dependence of the X-ray scattered intensity $I(q, \phi = \theta + \frac{\pi}{2})$ on the momentum transfer, q , for two values of the angle θ . The curves have been offset vertically for clarity, otherwise the data points would coincide at high q . Ref.[5]

The coefficient A in (13), (14) is found from normalization condition Ref.[1]. It is significant that the structure factor can be directly measured in the small angle X-ray scattering measurements. The results of such measurements for an uniaxially deformed aerogel are represented in Ref.[5]. As it was shown by the authors of Ref.[5] the results of their measurements can be fitted if the assumption is made that the fractal dimension of aerogel is not changed by the deformation and the correlation radius gains the following angular dependence (fig.2):

$$R(\theta) = R_0 + r_1 \cos(2\theta), \quad (15)$$

where R_0 - the correlation radius before deformation, θ - the angle between deformation axis and chosen direction. Amplitude of changing of correlation radius can be expressed via macroscopic parameters characterizing strain of aerogel:

$$\frac{r_1}{R_0} = \frac{1}{2}(1 - \nu)\alpha \frac{\Delta l}{l_0}, \quad (16)$$

here $\frac{\Delta l}{l_0}$ relative change of aerogel length, ν - the macroscopic Poisson ratio, ($\nu \approx 0.2 \div 0.3$), α - the coefficient of transmission of macroscopic strain down to the level of correlation length. It was found that α approximately equals to 3.

Substitution of the new structure factor into (10) yields the anisotropic

correction to the Self-energy part. Its principal order contribution arises from angular dependence of the correlation radius and k-dependence of the unperturbed Green's function. For the anisotropic part of the Self-energy function in the second order correction one arrives at:

$$\Sigma_{jl}^{(2)a} \approx 0.35 \left(\frac{\pi}{3}\right)^4 \frac{3-D}{D-1} \frac{R_0 \cdot r_1}{l_{tr}^2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}. \quad (17)$$

The third order correction has the form:

$$\Sigma_{jl}^{(3)} = n \left(\eta^{(1)}(0)\right)^3 \int \frac{d^3 k_1}{(2\pi)^3} \frac{d^3 k_2}{(2\pi)^3} G_{jm}^{(0)}(\mathbf{k}_1) G_{lm}^{(0)}(\mathbf{k}_2) \langle \sum_{t,u} e^{i\mathbf{k}_1 \mathbf{r}_{st}} e^{i\mathbf{k}_2 \mathbf{r}_{su}} \rangle \quad (18)$$

According to Ref.[1] it can be calculated with the assumption that:

$$\langle \sum_{t,u} e^{i\mathbf{k}_1 \mathbf{r}_{st}} e^{i\mathbf{k}_2 \mathbf{r}_{us}} \rangle = S(\mathbf{k}_1) S(\mathbf{k}_2). \quad (19)$$

If the same assumption is made for the high order corrections, sum of series, which forms geometric progression, can be found. In particular for the anisotropic part of the Self-energy one has:

$$\tau_{jl}^a = \Sigma_{jl}^a \approx 0.35 \left(\frac{\pi}{3}\right)^4 \frac{3-D}{D-1} \frac{R \cdot r_1}{l_{tr}^2} \frac{1}{\left(1 + \frac{25}{27} \frac{\pi^2}{2} \frac{R^2}{\xi_0 l_{tr}} \frac{1}{D-1}\right)^2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}. \quad (20)$$

As it is seen from the final formula in the case when $\xi_0 \gg R$ HSM can be used. In the opposite case correlations play an important role in the orientational effect.

Let us first consider the orientational effect of the anisotropic aerogel on the order parameter of the B-like phase. In the absence of magnetic field the order parameter of the B-like phase is isotropic and no orientational effect is expected. But situation changes in the presence of a magnetic field. The order parameter of the B-like phase in magnetic field has the form [6]:

$$A_{\mu j} = \frac{e^{i\phi}}{\sqrt{3}} \left(\Delta_{\perp} R_{\mu j} + (\Delta_{\parallel} - \Delta_{\perp}) R_{\nu j} \hat{h}_{\nu} \hat{h}_{\mu} \right), \quad (21)$$

the rotation matrix $R_{\mu j}(\hat{\mathbf{n}}, \theta)$ is characterized by a rotation axis $\hat{\mathbf{n}}$ and the rotation angle $\theta_L = \arccos(-\frac{1}{4} \frac{\Delta_{\parallel}}{\Delta_{\perp}})$, which is found from minimization of dipole energy, $\hat{\mathbf{h}}$ is the direction of magnetic field. Orientation of $\hat{\mathbf{n}}$ is




Figure 3:

Changing in orientation of vectors $\hat{\mathbf{n}}$ and $\hat{\mathbf{l}}$ in the presence of uniaxially-stretched aerogel.

induced by magnetic field and dipole forces. The corresponding term in the free energy equals to:

$$\Delta F_H^B = \lambda_D N_F (\Delta_{\parallel} - \Delta_{\perp}) \Delta_{\perp} (\hat{n} \cdot \hat{h})^2. \quad (22)$$

At the same time globally anisotropic aerogel induces orientation of the orbital part of the order parameter. Substitution of (21) into (4) gives:

$$\Delta F_a^B = \frac{1}{3} N_F (\Delta_{\parallel}^2 - \Delta_{\perp}^2) R_{\mu j} R_{\nu l} h_{\mu} h_{\nu} \tau_{jl}^a \approx \frac{2}{3} (\Delta_{\parallel} - \Delta_{\perp}) \Delta_{\perp} \Sigma_{xx}^a (1 - 3l_z^2), \quad (23)$$

here the orbital vector

$$l_j = \hat{R}_{\mu j} S_{\mu}, \quad (24)$$

is introduced, S_{μ} - the spin vector. Let us compare orientational energies for the case of uniaxially stretched aerogel in the direction of the magnetic field. In the absence of anisotropy vectors $\hat{\mathbf{n}}$, $\hat{\mathbf{l}}$ and $\hat{\mathbf{s}}$ are directed along the magnetic field fig.3. When aerogel is stretched vector $\hat{\mathbf{l}}$ tends to turn in the direction perpendicular to the magnetic field. To realize this it is needed to turn vector $\hat{\mathbf{n}}$. Taking into account the minimum of dipole energy solutions of Eq. (24) for a given configuration of vectors $\hat{\mathbf{n}}$, $\hat{\mathbf{l}}$ and $\hat{\mathbf{s}}$ always exist. The new configuration starts to form when:

$$\Delta F_a^B \sim \Delta F_H^B. \quad (25)$$

Estimations made for the whole interval of parameters characterizing aerogel and superfluid ^3He , $D = 1.6 \div 1.9$, $R \sim 200 \div 1400 \text{ \AA}$, $\xi_0 \sim 170 \div 300 \text{ \AA}$, $l_{tr} \approx 1400 \text{ \AA}$ show that deformation on the order of

$$\gamma_{min} \sim 5 \cdot (10^{-3} \div 10^{-4}) \quad (26)$$

is already strong enough to achieve orientation of the order parameter which is different from that required by the magnetic field in bulk liquid.

Estimation of the strength of the orientational effect in the A-like phase can be made in a similar way. For clarity we assume that aerogel is uniaxially squeezed. As it was shown in Ref.[7], [8] the order parameter of the A-like phase in the case of uniaxially-squeezed aerogel has the form of the bulk A-phase. Consequently the order parameter of the A-like phase can be written in the form:

$$A_{\mu j} = \frac{1}{\sqrt{2}} d_{\mu} (m_j + i n_j), \quad (27)$$

where vector $\hat{\mathbf{d}}$ corresponds to the spin part of the order parameter, and vector $\mathbf{l} = [\mathbf{m}, \mathbf{n}]$ is its orbital part. In the absence of squeezed aerogel magnetic field and dipole forces orient vector $\hat{\mathbf{l}}$ perpendicular to the magnetic field:

$$\Delta F_H^A = \frac{1}{2} \Delta \chi (\hat{\mathbf{d}} \cdot \mathbf{H})^2, \quad (28)$$

$$\Delta f_D = -\frac{2}{5} \lambda_D N_F \Delta_0^2 (\hat{\mathbf{d}} \cdot \hat{\mathbf{l}})^2. \quad (29)$$

However, as in the case $^3\text{He} - B$ aerogel can orient the orbital part of the order parameter:

$$\Delta F_a^A = -N_F \frac{\Delta^2}{2} \Sigma_{xx}^a (1 - 3l_z^2). \quad (30)$$

Comparison of ΔF_a^A with ΔF_H^A can be used for estimation of the strength of the orientational effect. Calculations show that for the melting pressure magnetic field corresponding to one percent squeezing approximately equals to 80G.

Comparison of Eqns. (23), (30) with the analogous formula from Ref.[2] shows that correlations can significantly decrease effect of anisotropy (on the order of 10). Let us note also that the effect of correlations on orientation of the order parameter is stronger than the similar effect on the transition temperature.

I am grateful to I.A. Fomin for fruitful discussion. This work is partly supported by RFBR grant (no. 07-02-00214), by Ministry of Education and Science of Russian Federation, Russian Science-Support Fund and Landau Scholarship (A.F.) from Forschungszentrum Jülich, Germany

References

- [1] I.A. Fomin, arXiv:cond-mat/0806.1469v1

- [2] E.V. Surovtsev, I.A. Fomin, J Low Temp Phys **150** (2008), Numbers 3-4, 487-492
- [3] D. Rainer, M. Vuorio J.Phys. C: Solid State Phys. **10** (1977) 3093.
- [4] I.A. Fomin, J. of Phys. and Chemistry of Solids **66** (2005) 1321-1324.
- [5] J. Pollanen, K. Shirer, S. Blinstein, J.P. Davis, H. Choi, T.M. Lippman, W.P. Halperin, arXiv:cond-mat/0711.3495v1
- [6] D. Vollhardt, P. Wölfle, The Superfluid Phases of 3He , Taylor and Francis, (1990), 134-135
- [7] I.A. Fomin, J Low Temp Phys **150** (2008), Numbers 3-4, 464-471
- [8] G.E. Volovik, J Low Temp Phys **150** (2008), Numbers 3-4, 453-463